IN THE CLAIMS

The following listing of claims will replace all prior versions, and listings, of claims in this application.

Listing of the Claims:

Claims 1-21 (Cancelled).

22. (Original) An isolated polypeptide comprising SEQ ID NO:2.

Claims 23-25 (Cancelled).

26. (Currently Amended) An N-acetylglycosamine-1-phosphotransferase (GlcNAc-phosphotransferase) comprising an α subunit, a β subunit and a site-specific proteolytic cleavage site interposed between said α and β subunits, wherein said site-specific proteolytic cleavage site is not endogenous to GlcNAc-phosphotransferase,

wherein said α subunit is encoded by nucleotides 165 to 2948 of SEQ ID NO:3, or a sequence that hybridizes under stringent conditions to the complement of nucleotides 165 to 2948 of SEQ ID NO:3, wherein said stringent conditions comprise hybridization in 50% formamide, 1M NaCl, 1% SDS at 37°C and washing in 0.1 X SSC at 60 to 65°C and which encodes a protein when combined with a β subunit protein encoded by nucleotides 2949 to 2932 of SEQ ID NO:3 has GlcNAc-phosphotransferase activity; and

wherein said β -subunit is encoded by nucleotides 2949 to 3932 of SEQ ID NO:3, or a sequence that hybridizes under stringent conditions to the complement of nucleotides 2949 to 3932 of SEQ ID NO:3, wherein said stringent conditions comprise hybridization in 50% formamide, 1M NaCl, 1% SDS at 37°C and washing in 0.1 X SSC at 60-to 65°C and which encodes a protein when combined with an α subunit protein encoded by nucleotides 165 to 2948 of SEQ ID NO:3 has GlcNAc-phosphotransferase activity.

Claims 27-29 (Cancelled).

- 30. (Previously Presented) The GlcNAc-phosphotransferase of Claim 26, wherein said α -subunit comprises amino acids 1-928 of SEQ ID NO:4.
- 31. (Previously Presented) The GlcNAc-phosphotransferase of Claim 26, wherein said β subunit comprises amino acids 1 to 328 of SEQ ID NO:5.
- 32. (Original) The GlcNAc-phosphotransferase of Claim 26, wherein said GlcNAc-phosphotransferase further comprises a γ subunit.

Claim 33 (Cancelled).

- 34. (Original) The GlcNAc-phosphotransferase of Claim 32, wherein said γ subunit comprises the amino acid sequence of SEQ ID NO:7.
- 35. (Currently Amended) The GlcNAc-phosphotransferase of Claim 26, wherein said site-specific proteolytic cleavage site is selected from the group consisting of a Furin proteolytic cleavage site, a Factor Xa proteolytic cleavage site, a Enterokinase proteolytic cleavage site, and a Genenase I proteolytic cleavage site comprising SEQ ID NO:25.
- 36. (Original) The GlcNAc-phosphotransferase of Claim 35, wherein said site-specific proteolytic cleavage site is a Furin proteolytic cleavage site.
- 37. (Previously Presented) The GlcNAc-phosphotransferase of Claim 36, wherein said Furin proteolytic cleavage site comprises SEQ ID NO:24.

Claims 38-55 (Cancelled).

56. (Previously Presented) A method of phosphorylating a lysosomal hydrolase protein comprising an asparagine-linked oligosaccharide with a high mannose structure, the method comprising contacting said lysosomal hydrolase protein with the isolated polypeptide of Claim 22 for a time and under conditions suitable to produce a phosphorylated lysosomal hydrolase protein.

Claims 57-69 (Cancelled).

- 70. (Currently Amended) The method of Claim 56, wherein said lysosomal hydrolase protein is selected from the group consisting of α -glucosidase, α -L-iduronidase, α -galactosidase A, arylsulfatase, N-acetlygalactosamine- α -sulfatase, β -galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, β -glucoronidase, Heparan N-sulfatase, N-Acetyl- α -glucosaminidase, Acetyl CoA- α glucosaminide N acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside sialidase, Acid β -galactosidase G_{M1} -Ganglioside, Acid- β -galactosidase, Hexosaminidase A, Hexosaminidase B, α -fucosidase, α -N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β -Glucosidase.
- 71. (Previously Presented) The method of Claim 56, further comprising contacting said phosphoryalated protein with an isolated N-acetylglucosamine-1-phosphodiester-N-Acetylglucosaminidase (phosphodiester α -GlcNAcase).
- 72. (Previously Presented) The method of Claim 71, wherein said phosphodiester α -GlcNAcase comprises the amino acid sequence of SEQ ID NO:18.
- 73. (Currently Amended) The method of Claim 71, wherein said phosphodiester α -GlcNAcase is encoded by a nucleotide sequence comprising SEQ ID NO:17 or a nucleotide sequence that hybridizes under stringent conditions to the complement of SEQ ID NO:17, wherein said stringent conditions comprise hybridization in 50% formamide, 1M NaCl, 1% SDS at 37°C and washing in 0.1 X SSC at 60 to 65°C, and which encodes a protein with phosphodiester α -GlcNAcase activity.
- 74. (Previously Presented) The method of Claim 56, wherein prior to said contacting the method comprises: culturing a host cell which comprises a polynucleotide encoding the

polypeptide for a time under conditions suitable for expression of the polypeptide; and isolating said polypeptide.

75. (Cancelled)

76. (Currently Amended) A method of phosphorylating a lysosomal hydrolase protein comprising an asparagines asparagine-linked oligosaccharide with a high mannose structure, the method comprising contacting said protein with the GlcNAc phosphotransferase of Claim 26 for a time and under conditions suitable to produce a phosphorylated protein.

Claims 77-80 (Cancelled).

- 81. (Previously Presented) The method of Claim 7678, wherein said α -subunit comprises amino acids 1-928 of SEQ ID NO:4.
- 82. (Previously Presented) The method of Claim 76 78, wherein said β subunit comprises amino acids 1 to 328 of SEQ ID NO:5.
- 83. (Previously Presented) The method of Claim 76 78, wherein said soluble GlcNAcphosphotransferase further comprises a γ subunit.
- 84. (Currently Amended) The method of Claim 83, wherein said γ subunit is encoded by SEQ ID NO:6, or a nucleotide sequence that hybridizes under stringent conditions to the complement of SEQ ID NO:6, wherein said stringent conditions comprise hybridization in 50% formamide, 1M NaCl, 1% SDS at 37°C and washing in 0.1 X SSC at 60 to 65°C.
- 85. (Previously Presented) The method of Claim 83, wherein said γ subunit comprises the amino acid sequence of SEQ ID NO:7.
- 86. (Currently Amended) The method of Claim 76, wherein said site-specific proteolytic cleavage site is selected from the group consisting of a Furin proteolytic cleavage

site, a Factor Xa proteolytic cleavage site, a Enterokinase proteolytic cleavage site, and a Genenase I-proteolytic cleavage site comprising SEQ ID NO:25.

- 87. (Previously Presented) The method of Claim 86, wherein said site-specific proteolytic cleavage site is a Furin proteolytic cleavage site.
- 88. (Previously Presented) The method of Claim 87, wherein said Furin proteolytic cleavage site comprises SEQ ID NO:24.

Claim 89 (Cancelled).

- 90. (Currently Amended) The method of Claim 76 89, wherein said lysosomal hydrolase protein is selected from the group consisting of α-glucosidase, α-L-iduronidase, α-galactosidase A, arylsulfatase, N-acetlygalactosamine-α -sulfatase, β-galactosidase, iduronate 2-sulfatase, ceramidase, galactocerebrosidase, β-glucoronidase, Heparan N-sulfatase, N-Acetyl-α-glucosaminidase, Acetyl CoA α-glucosaminide N-acetyl transferase, N-acetyl-glucosamine-6 sulfatase, Galactose 6-sulfatase, Arylsulfatase A, Arylsulfatase B, Arylsulfatase C, Arylsulfatase A Cerebroside, Ganglioside sialidase, Acid β-galactosidase G_{MI}-Ganglioside, Acid β-galactosidase, Hexosaminidase A, Hexosaminidase B, α-fucosidase, α-N-Acetyl galactosaminidase, Glycoprotein Neuraminidase, Aspartylglucosamine amidase, Acid Lipase, Acid Ceramidase, Lysosomal Sphingomyelinase, Sphingomyelinase, and Glucocerebrosidase β-Glucosidase.
- 91. (Previously Presented) The method of Claim 76, further comprising contacting said phosphoryalated protein with an isolated phosphodiester α -GlcNAcase.
- 92. (Previously Presented) The method of Claim 91, wherein said phosphodiester α -GlcNAcase comprises the amino acid sequence of SEQ ID NO:18.
- 93. (Currently Amended) The method of Claim 91, wherein said phosphodiester α -GlcNAcase is encoded by a nucleotide sequence comprising SEQ ID NO:17 or a nucleotide

sequence that hybridizes under stringent conditions to the complement of SEQ ID NO:17, wherein said stringent conditions comprise hybridization in 50% formamide, 1M NaCl, 1% SDS at 37°C and washing in 0.1 X SSC at 60 to 65°C.

94. (Previously Presented) The method of Claim 76, wherein prior to said contacting the method comprises: culturing a host cell which comprises a polynucleotide encoding the GlcNAc-phosphotransferase for a time under conditions suitable for expression of the GlcNAc-phosphotransferase; and isolating said GlcNAc-phosphotransferase.

95. (Cancelled)